

# Vitamine E

## Journal of Cosmetic Science

52, 155-161 (May/June 2001)

Direct evidence for bioconversion of vitamin E acetate into vitamin E:  
An ex vivo study in viable human skin

*W. BASCHONG, C. ARTMANN, D. HUEGLIN, and J. ROEDING, M. E. Mueller Institute at the Biozentrum, University of Basel, Klingelbergstrasse 50-70, CH-4056 Basel, Switzerland (W.B.), PhaCos GmbH, Munich, Germany (C.A.), Ciba Specialty Chemicals, P.O. Box 1266, D-79630 Grenzach-Wyhlen, Germany (D.H.), and Mica Products GmbH, Zoellinplatz 4, D-79410 Badenweiler, Germany (J.R.). Accepted for publication March 15, 2001.*

### *Synopsis*

For better stability, vitamin E is commonly used as the non-active esterified pro-drug. Such esters are postulated to be hydrolyzed to the free active form by skin-related esterases. So far, successful conversion of esterified vitamin E to free vitamin E (tocopherol) has been mainly delineated from observed biological effects. Quantitative evidence in human skin is poor. In vitro and in vivo studies on human and animal skin have proved ambiguous. Formulation-based effects may have added to this controversy. **In the present study, comparable amounts of vitamin E acetate (i) in oil (Mygliol-812N), (ii) surfactantsolubilized in water, (iii) encapsulated in liposomes, or (iv) encapsulated in Nanotopes™ were applied to human skin** mounted in modified Franz-perfusion chambers that permit emulation of both open or occlusive conditions. The distribution of vitamin E total (vitamin E acetate + vitamin E) was assessed on the skin surface, in the horny layers, and in the underlying skin by high-pressure liquid chromatography (HPLC), with a recovery higher than 90%. **Vitamin E acetate in Mygliol deposited exclusively on the surface and in the stratum corneum. In contrast, solubilized or encapsulated vitamin E acetate deposited also in the underlying skin. Nanotopes™ performed best, followed by liposomes and solubilized vitamin E acetate.** Non-occlusive application favored deposition in the skin relative to occlusive application. **Conversion of vitamin E acetate to vitamin E was not observed on the skin surface or in the horny layers, while in the underlying skin up to 50% of the vitamin E total was deacetylated.**